arXiv:1210.1452v1 [cond-mat.stat-mech] 4 Oct 2012

Casimir force in the $O(n \to \infty)$ model with free boundary conditions

Jonathan Bergknoff[1], Daniel Dantchev[1,2] and Joseph Rudnick[1]
[1] *Department of Physics and Astronomy, UCLA, Los Angeles, California 90095-1547, USA,*
[2] *Institute of Mechanics - BAS, Academic Georgy Bonchev St. building 4, 1113 Sofia, Bulgaria*

(Dated: October 5, 2012)

We study the Casimir force in a film with free boundary conditions described by the $n \to \infty$ limit of the $O(n)$ models. Via exact calculations we reduce the model to an one dimensional model the properties of which we determine numerically, near the critical region, and in an exact analytical manner in the low-temperature regime. The results resemble the experimental curve behavior for the force obtained for ^{4}He films.

PACS numbers: 64.60.-i, 64.60.Fr, 75.40.-s

I. INTRODUCTION

A recent paper by Diehl *et. al.* [1] reports on a study of the Casimir force in thin films (i.e. dimensions $\infty^2 \times L$) of a Ginzburg-Landau-Wilson (GLW) version of the $O(n)$ model in the limit $n \to \infty$, the system being subject to free boundary conditions in the finite direction. Because of the continuous symmetry of the model, which is broken at low temperature when $L \to \infty$, as well as the fact that the boundary conditions correspond to those that are appropriate in the case of ^{4}He films the results of the calculations in [1] are relevant to the Casimir force measurements on such films described in [2, 3]. The superfluid transition in ^{4}He is, of course, correctly modeled in terms of the XY, or $O(2)$, model, and the results in refs. [2, 3] have been quite successfully reproduced by Monte Carlo simulations of this model in [4] and [5].

Nevertheless, the $O(n \to \infty)$ model merits consideration as a depiction of systems with broken continuous symmetry in the bulk insofar as this model is susceptible to a combination of analytical and numerical approaches, yielding both quantitative and qualitative insights into the behavior of those systems. In the current article we study the $n \to \infty$ limit of the $O(n)$ models without using the mapping on the GLW model, but relying on the definition of the corresponding spherical model [6]; we recall that the infinite translational invariant standard spherical model is equivalent to the $n \to \infty$ limit of the corresponding system of n-component vectors [7, 8], but for the spherical model with surfaces or, more generally, without translation-invariant symmetry, this equivalence is preserved only if one imposes spherical constraints in a way which ensures that the mean square value of each spin of the system is the same [9]—that is, one averages thermally, but not spatially.

Generally such a model is considered analytically intractable. However, as we demonstrate here, this model can be analytically reduced via exact calculations to an one dimensional model the properties of which can be then either studied numerically, near the critical region, or in an exact analytical manner in the low-temperature regime. The differences of our approach from the one presented in Ref. [1] are *i)* the model is self-contained and does not require the determination of any parameters, e.g. the strength of the fourth order coupling in the GLW model, and *ii)* there is no need to use an effective width of the system, keeping in mind that there is always some small (microscopic) ambiguity in the definition of the thickness L; see, e.g., Ref. [10] and the discussion therein. Finally, the present formulation of the model is, unlike the GLW approach, suitable for investigation of the properties of the system at all temperatures, particularly at temperatures considerably below that of the bulk transition and thus well outside of the critical regime.

It is worth reiterating that the Casimir effect remains the object of intense studies, in its original formulation due to Casimir [11] (see the reviews [12, 13]), and in its thermodynamic manifestation [14]—see, e.g, the general reviews [15, 16] and the reviews devoted to some specific aspects of the critical Casimir force [17, 18].

II. DEFINITION OF THE MODEL

For an $O(n), n \geq 1$ model of a d-dimensional system at a temperature T and geometry $\infty^{d-1} \times L$ the thermodynamic Casimir force is defined by [19], [15]

$$F^{(\tau)}_{\text{Casimir}}(T, L) = -\frac{\partial f^{(\tau)}_{\text{ex}}(T, L)}{\partial L}, \tag{2.1}$$

where $f^{(\tau)}_{\text{ex}}(T, L)$ is the excess free energy

$$f^{(\tau)}_{\text{ex}}(T, L) = f^{(\tau)}(T, L) - L f_b(T), \tag{2.2}$$

and the superscript τ denotes the boundary conditions. Here $f^{(\tau)}(T, L)$ is the full free energy per unit area of such a system subjected to the boundary conditions τ and f_b is the bulk free energy density.

Consider a d-dimensional square lattice, each lattice site occupied by an n-component classical vector spin having ferromagnetic interactions with its nearest neighbors. We single out one dimension, z, to be L lattice spacings long. At each of the L sites along the finite dimension, there is a $(d-1)$-dimensional transverse layer containing a total of A spins, where A is large and will later be taken to infinity. Free boundary conditions are imposed in the z direction by putting a layer of zero length spins on the top and the bottom of the film (i.e. at $z = 0$ and $z = L + 1$).

The model, as described, is not especially amenable to analysis. However, in the $n \to \infty$ limit it is equivalent[7, 9] to a form of the spherical model, wherein the vector spins are replaced by real-valued scalar spins and each $(d-1)$-dimensional layer satisfies an individual spherical constraint $\sum s^2 = A$. We simplify matters further by using the mean spherical model, which gives the same results as the spherical model in the thermodynamic limit, i.e. with $A \to \infty$.

Our Hamiltonian is therefore

$$H = -J \sum_{\langle s,s' \rangle} ss' + J \sum_i \Lambda_i \left(\sum_j s_{i,j}^2 - A \right) \tag{2.3}$$

where the first summation is taken over nearest neighbor spins s and s', which may each lie in the same layer or in adjacent layers. $J > 0$ is the ferromagnetic coupling and Λ_i is the spherical field for layer i which will be used to enforce the mean spherical constraints. The notation $s_{i,j}$ refers to spin j in layer i, with $i = 1, \ldots, L$ and $j = 1, \ldots, A$.

Fourier transforming spins according to

$$s_{i,j} = A^{-1/2} \sum_{\mathbf{q}} s_i(\mathbf{q}) e^{i \mathbf{r}_j \cdot \mathbf{q}}, \tag{2.4}$$

where the sum extends over the first Brillouin zone of layer i, we find

$$H = -JA \sum_i \Lambda_i + \frac{J}{2} \sum_{\mathbf{q}} \mathbf{s}(\mathbf{q})^\dagger \mathcal{H}(\mathbf{q}) \mathbf{s}(\mathbf{q}) \tag{2.5}$$

with

$$[\mathcal{H}(\mathbf{q})]_{ij} = \mathcal{M}_{ij} - 2\delta_{ij} \sum_{k=1}^{d-1} \cos q_k \tag{2.6}$$

and

$$\mathcal{M}_{ij} = \begin{cases} 2\Lambda_i, & i = j \\ -1, & |i-j| = 1 \\ 0, & \text{otherwise} \end{cases}. \tag{2.7}$$

After computing the partition function in the standard way, we find that the free energy per transverse unit area, in units of $k_B T$, to be

$$\frac{\beta \mathcal{F}}{A} = -\beta J \sum_i \Lambda_i + \frac{1}{2} L \ln \left(\frac{\beta J}{2\pi} \right) + \frac{1}{2A} \sum_{\mathbf{q}} \ln \left[\det \left(\mathcal{H}(\mathbf{q}) \right) \right]. \tag{2.8}$$

The spherical constraint is enforced in the mean via the Lagrange multipliers Λ_i. In particular, we must have

$$0 = \frac{\beta}{A} \frac{\partial \mathcal{F}}{\partial \Lambda_i} = -\beta J + \frac{1}{A} \sum_{\mathbf{q}} \left[\mathcal{H}(\mathbf{q}) \right]_{ii}^{-1} \tag{2.9}$$

for each $i = 1, \ldots, L$.

III. RESULTS ON THE MODEL IN $d = 3$

We specialize now to the case of three dimensions. We write $\mathcal{H}^{-1}$ using an outer-product expansion. Taking $\{a_l, |\psi^{(l)}\rangle\}$ to be the eigenvalues and normalized eigenvectors, respectively, of the matrix $\mathcal{M}$ defined in Eqn. (2.7), we have

$$\mathcal{H}(\mathbf{q})^{-1} = \sum_{l=1}^{L} \frac{|\psi^{(l)}\rangle \langle \psi^{(l)}|}{a_l - 2\cos q_x - 2\cos q_y}. \tag{3.1}$$

In the transverse thermodynamic limit, $A \to \infty$, the sum over Brillouin zone is reproduced by an integral, i.e. $\sum_{\mathbf{q}} \to \frac{A}{(2\pi)^2} \int d^2 q$. In light of that, Eqns. (2.9) and (3.1) give three-dimensional spherical constraints

$$\beta J = \frac{1}{2\pi} \sum_{l=1}^{L} \psi_i^{(l)\,2} \, \frac{4}{a_l} K \left(\frac{4}{a_l} \right), \tag{3.2}$$

for each $i = 1, \ldots, L$, where $K(k)$ is the complete elliptic integral of first kind with modulus k.

Furthermore, the free energy can be written in closed form for $d = 3$:

$$\frac{\beta \mathcal{F}}{A} = \frac{1}{2} L \ln \left(\frac{\beta J}{2\pi} \right) + \frac{1}{2} \sum_{l=1}^{L} \Bigg[\ln a_l - 2\beta J \Lambda_l - \frac{2}{a_l^2} \cdot {}_4F_3 \left(1, 1, \frac{3}{2}, \frac{3}{2}; 2, 2, 2; \frac{16}{a_l^2} \right) \Bigg], \tag{3.3}$$

with generalized hypergeometric function ${}_4F_3$. The bulk free energy per length, f_b, is also computed to be

$$\beta f_b = -\beta J \Lambda + \frac{1}{2} \ln \left(\frac{\beta J}{2\pi} \right) + \frac{1}{4\pi^2} \int_{-\pi}^{\pi} dq_x \, dq_y \, \sinh^{-1} u, \tag{3.4}$$

with $u = \frac{1}{2}\sqrt{4 - 2\cos q_x - 2\cos q_y}$ and $\Lambda = 3$ for $\beta J > \beta_c J$, while Λ for $\beta J \le \beta_c J$ is determined by the bulk constraint

$$\beta J = \int_{-\pi}^{\pi} \frac{d^3 q}{\Lambda - \cos q_x - \cos q_y - \cos q_z}, \tag{3.5}$$

where bulk critical coupling $\beta_c J$ is provided by the above equation via setting $\Lambda = 3$ in it.

Therefore, given a temperature, the problem is to solve Eqs. (3.2) simultaneously, with $i = 1, \ldots, L$, for the unknown spherical fields, Λ_i. Then, we construct the Casimir pressure

$$\frac{\beta F_{\text{Cas}}}{A} = -\frac{\partial}{\partial L} \left(\frac{\beta \mathcal{F}}{A} - L \beta f_b \right) \approx \beta f_b - \frac{1}{2} \left[\frac{\beta \mathcal{F}(L+1)}{A} - \frac{\beta \mathcal{F}(L-1)}{A} \right]. \tag{3.6}$$

Implementing this analysis numerically, we find the Casimir force curves shown in Fig. 1 for several system sizes L. The scaling variable is $x = (L/\xi_0^+)t$, where $t = (T - T_c)/T_c$ is the reduced temperature and $\xi_b(t \to 0^+) = \xi_0^+ t^{-\nu}$ is the bulk correlation length. Here, $\nu = 1$ is the corresponding critical exponent in the bulk spherical model in $d = 3$, and $\xi_0^+ = [4\pi(\beta_c J)]^{-3/2}$ the non-universal amplitude as determined from earlier results [20]. As $L \to \infty$, the scaling corrections vanish and we are left with the Casimir scaling function $\vartheta(x) = L^3 \beta F_{\text{Cas}}(x)/A$ for this system under free boundary conditions. In fact, the curves for $L = 200$ and $L = 500$ are substantially the same, so the solid black $L = 500$ curve is already an excellent approximation to $\vartheta(x)$.

We note here that the full range of negative x appropriate to each curve depends on L; the $T = 0$ limit is achieved at $x = -L/\xi_0^+$.

From our numerical results, we find a Casimir amplitude of

$$\Delta = \frac{1}{2}\vartheta(0) \approx -0.0107737 \tag{3.7}$$

and an extremum of the scaling function

$$\vartheta_{\min} \approx -0.1269001. \tag{3.8}$$

These quantities are expected to be universal and, indeed, they agree to great precision with the values given by Diehl, *et. al.* [1] for their closely related model. The location of the Casimir force extremum is at $x = x = -8.08$ for $L = 500$. Agreement with the measurements in [2, 3] is less satisfactory, as it should be given the widely-acknowledged difference between the $O(2)$ and $O(\infty)$ models.



FIG. 1. (Color online) The scaled Casimir force $L^3 \beta F_{\text{Cas}}/A$ as a function of scaling variable $x = (L/\xi_0^+)t$, with reduced temperature $t = (T - T_c)/T_c$ and bulk correlation length amplitude ξ_0^+, for $L = 10$ (top, solid blue), $L = 30$ (second from top, dashed purple), $L = 200$ (second from bottom, dot-dashed red), $L = 500$ (bottom, solid black). The zero-temperature value of $-\zeta(3)/8\pi$ is indicated as a horizontal dashed line. The $L = 200$ and $L = 500$ curves lie on top of eachother in the critical region $x \approx 0$, and both are close approximations to the scaling function $\vartheta(x)$.

IV. ASYMPTOTIC LOW TEMPERATURE RESULTS IN $d = 3$

The system may be studied in closed form for $T \ll T_c$, i.e. $\beta J \gg \beta_c J$. In that regime, the left hand side of Eqn. (3.2) becomes large, forcing the lowest eigenvalue, a_1, to approach 4 and dominate the right hand side. This effectively renders the constraint as

$$\beta J \approx \frac{1}{2\pi}\, \psi_i^{(1)\,2}\, \frac{4}{a_1} K\left(\frac{4}{a_1}\right). \tag{4.1}$$

As this must hold for each i, we will have $\psi_i^{(1)\,2} = \text{const.}$ Numerically, we see that this eigenvector approaches a constant vector (as opposed to one which varies with respect to i). In order for the constant vector to be an eigenvector of $\mathcal{M}$ with eigenvalue 4, we must have $\mathbf{\Lambda} = \mathbf{\Lambda}^* = (5, 6, 6, \ldots, 6, 6, 5)/2$. We now expand about $\mathbf{\Lambda} = \mathbf{\Lambda}^*$ using perturbation theory.

The eigenvalues and normalized eigenvectors of $\mathcal{M}(\mathbf{\Lambda}^*)$ may be found in closed form as

$$\lambda_l = 4 + 4\sin^2\left(\frac{\pi(l-1)}{2L}\right) \tag{4.2}$$

and

$$\phi_i^{(l)} = \cos\left(\frac{\pi(l-1)}{2L}(2i-1)\right)/\text{norm}, \tag{4.3}$$

where $l = 1, \ldots, L$ and the norm is $\sqrt{L}$ for $l = 1$ and $\sqrt{L/2}$ for all other l. Applying perturbation theory and using the constraint, Eqn. (3.2), we find

$$a_1 \approx 4 + 32 e^{-4\pi L(R - R_c)}, \tag{4.4}$$

while

$$\begin{aligned} a_l = \lambda_l &+ \frac{1}{2\pi L(R - R_c)}(\lambda_{2l-1} - 4) \\ &\times \left[\frac{K(4/\lambda_l)}{\lambda_l} - \frac{K(4/\lambda_{L+2-l})}{\lambda_{L+2-l}}\right] + O\left(R^{-2}\right), \end{aligned} \tag{4.5}$$

for $l \geq 2$, where we now abbreviate $R = \beta J$ and R_c the corresponding value at the bulk critical point. As before, $K(k)$ refers to the elliptic integral with modulus k, and λ_n is given by Eqn. (4.2) even when n is not in the range $1, \ldots, L$. Note that there is one eigenvalue, a_1, which exhibits markedly different behavior from the other eigenvalues.

In order to compute the Casimir force, we must determine the effects of the perturbed eigenvalues on the free energy, Eqns. (3.3) and (3.4). A careful analysis, whose details are deferred to a later publication, shows



FIG. 2. (Color online) The scaled Casimir force (symbols) as compared to the closed form asymptotic results (solid curves) presented in section IV for $L = 50, 100, 200$ and 500. The asymptotic results turn out to be accurate for moderately low absolute temperatures, corresponding to reduced temperatures $t \leq -0.2$ in this plot. Note that the plot is versus *unscaled* reduced temperature, t, which is equal to -1 at $T = 0$.

that there is a leading order, temperature-independent, term in βF_{Cas}. This term is given by

$$-\frac{1}{16\pi^2}\int_{-\pi}^{\pi} d^2q \, \ln\left(\frac{1-e^{-4(L+1)v}}{1-e^{-4(L-1)v}}\right), \qquad (4.6)$$

where

$$v = \sinh^{-1}\left[\frac{1}{2}\sqrt{4-2\cos q_x - 2\cos q_y}\right]. \qquad (4.7)$$

Expanding in powers of $1/L$, we find

$$\frac{\beta F_{\mathrm{Cas}}}{A} = -\frac{1}{8\pi L^3}\left[\zeta(3) + \frac{2\zeta(3)+\frac{7}{8}\zeta(5)}{L^2} + O\left(L^{-4}\right)\right] + O(T). \qquad (4.8)$$

The leading temperature-dependent term is of the form

$$-\frac{1}{4(R-R_c)L^4}\left[a + b\ln L + O\left(L^{-2}\right)\right], \qquad (4.9)$$

with

$$a = \frac{\zeta'(-2)}{4}\left(2 + 3K\left(\frac{1}{2}\right) - 21\ln 2 + 6\ln(2\pi)\right) - \frac{3\zeta''(-2)}{4} \qquad (4.10)$$

and

$$b = -\frac{3\zeta'(-2)}{2}, \qquad (4.11)$$

where ζ is the Riemann ζ-function. Note that $1/(R - R_c) \sim T$ for $T \approx 0$. The comparison between numerical and asymptotic results is shown in Fig. 2, using in addition the next order results in powers of L which are omitted here for brevity.

While the Casimir forces in Figs. 1 and 2 differ in detail from the data for ^{4}He obtained in [2, 3], the overall features—particularly the pronounced minimum in the Casimir force below the critical point and the approach to a non-zero limiting value at low temperatures—are strikingly similar. This encourages us to anticipate that the model will prove to be a very useful adjunct to general, and perhaps specific, studies of Casimir forces in systems with a continuous symmetry of the type that is broken in the superfluid transition.

[1] H. W. Diehl, D. Grüneberg, M. Hasenbusch, A. Hucht, S. B. Rutkevich, and F. M. Schmidt, ArXiv e-prints (2012), 1205.6613.
[2] R. Garcia and M. H. W. Chan, Phys. Rev. Lett. **83**, 1187 (1999).
[3] A. Ganshin, S. Scheidemantel, R. Garcia, and M. H. W. Chan, Phys. Rev. Lett. **97**, 075301 (pages 4) (2006).
[4] A. Hucht, Phys. Rev. Lett. **99**, 185301 (2007).
[5] O. Vasilyev, A. Gambassi, A. Maciòłek, and S. Dietrich, Phys. Rev. E **79**, 041142 (2009).
[6] T. H. Berlin and M. Kac, Phys. Rev. **86**, 821 (1952).
[7] H. E. Stanley, Phys. Rev. **176**, 718 (1968).
[8] M. Kac and C. J. Thompson, Phys. Norveg. **5**, 163 (1971).
[9] H. J. F. Knops, Journal of Mathematical Physics **14**, 1918 (1973).
[10] D. Dantchev, F. Schlesener, and S. Dietrich, Phys. Rev. E **76**, 011121 (2007).
[11] H. B. Casimir, Proc. K. Ned. Akad. Wet. **51**, 793 (1948).
[12] R. Golestanian and M. Kardar, Phys. Rev. Lett. **78**, 3421 (1997).
[13] D. Dalvit, P. Milonni, D. Roberts, and F. E. da Rosa, eds., *Casimir Physics*, vol. 834 of *Lecture Notes in Physics* (Springer, Berlin, 2011), 1st ed.
[14] M. E. Fisher and P. G. de Gennes, C. R. Seances Acad. Sci. Paris Ser. B **287**, 207 (1978).
[15] J. G. Brankov, D. M. Dantchev, and N. S. Tonchev, *The Theory of Critical Phenomena in Finite-Size Systems -*

Scaling and Quantum Effects (World Scientific, Singapore, 2000).
[16] M. Krech, *Casimir Effect in Critical Systems* (World Scientific, Singapore, 1994).
[17] M. Krech, J. Phys.: Condens. Matter **11**, R391 (1999).
[18] A. Gambassi and S. Dietrich, Soft Matter **7**, 1247 (2011).
[19] R. Evans, *Liquids at interfaces* (Elsevier, Amsterdam, 1990).
[20] M. Barber and M. Fisher, Ann. of Phys. **77**, 1 (1973).